arXiv:1103.6171v1 [math.MG] 29 Mar 2011

Complexity of the Fibonacci snowflake

A. BLONDIN MASSÉ, S. BRLEK†, S.LABBÉ, AND M. MENDÈS FRANCE

A la mémoire de Benoit Mandelbrot

Abstract. The object under study is a particular closed curve on the square lattice $\mathbb{Z}^2$ related with the Fibonacci sequence F_n. It belongs to a class of curves whose length is $4F_{3n+1}$, and whose interiors by translation tile the plane. The limit object, when conveniently normalized, is a fractal line for which we compute first the fractal dimension, and then give a complexity measure.

§1. Introduction.

In a recent article [1] we defined and studied the properties of what we named a *Fibonacci snowflake*. Let us recall the main facts. Consider the infinite sequence $(q_n)_{n\in\mathbb{N}}$ defined recursively on the alphabet $\mathcal{T} = \{L = \text{left}, R = \text{right}\}$ by

$$\begin{aligned} q_0 &= \varepsilon \text{ (the empty word)}, \; q_1 = R, \\ q_n &= \begin{cases} q_{n-1}q_{n-2} & \text{if } n \equiv 2 \mod 3, \\ q_{n-1}\overline{q_{n-2}} & \text{if } n \equiv 0,1 \mod 3. \end{cases} \end{aligned}$$

Here, $\overline{q_{n-2}}$ represents the word q_{n-2} in which the letters R and L are permuted. The length $|q_n|$ of the word q_n satisfies the Fibonacci relation

$$|q_n| = |q_{n-1}| + |q_{n-2}|$$

and therefore

$$|q_n| = \frac{1}{\sqrt{5}}\left(\frac{1+\sqrt{5}}{2}\right)^n - \frac{1}{\sqrt{5}}\left(\frac{1-\sqrt{5}}{2}\right)^n.$$

To each word $w \in \mathcal{T}^* \cup \mathcal{T}^{\mathbb{N}}$ corresponds a polygonal line Π on the lattice $\mathbb{Z}^2$. At the n-th vertex of Π, the n-th letter w_n of w indicates the direction of the next side. It may well happen that some sides are visited several times as for the case $w = L^4$ for example. The length of

† With the support of NSERC (Canada).

the polygonal line Π corresponding to $w = L^4$ is : $|\Pi| = |L^4| + 1$. Actually, the following general property holds

$$|\Pi| = |w| + 1, \ \forall w \in \mathcal{T}^*. \tag{1}$$

A finite polygonal line on $\mathbb{Z}^2$ is *closed* if both extremities coincide. The corresponding word is then said to be closed. *Open* signifies non-closed; for example, L^4 is open while L^3 is closed. An open polygonal line is *non-intersecting* if each of its vertices is attained once only. A *closed non-intersecting* polygonal line is one for which the only vertices visited twice are the extremities. Here are a few examples:

(i.) L^4 is neither closed nor non-intersecting;
(ii.) L^3 is closed and non-intersecting;
(iii.) $(LR)^n$ is open and non-intersecting;
(iv) L^3RL^3 is closed but not non-intersecting;
(v) $(LRL)^3LR$ is closed and non-intersecting.

Given a word $w \neq \varepsilon$, we denote w^- the word where the last letter is suppressed. For example, $(LRL)^{4^-}$ is closed and non-intersecting. In [1] we showed that the polygonal line Π_n corresponding to the word $(q_{3n+1})^{4^-}$ is closed and non-intersecting (Theorem 10) (see Figure 1). Therefore, the property (1) implies that

$$|\Pi_n| = 4|q_{3n+1}|, \tag{2}$$

that is, the polygon Π_n is composed of $4|q_{3n+1}|$ unit segments in $\mathbb{Z}^2$.



Figure 1. Fibonacci snowflake of order $n = 5$.

§2. **Fractal dimension**. The rather "complicated" associated polygon Π_n, when conveniently normalized, tends to a fractal line as $n \to \infty$ which we call *Fibonacci snowflake*. In our previous article we were aware of its complexity, yet we overlooked its fractal dimension even though we had all the information needed. We now prove:

Theorem 1. *The fractal dimension of the Fibonacci snowflake is*

$$d = \frac{\log(2+\sqrt{5})}{\log(1+\sqrt{2})} = 1.637938210\cdots \tag{3}$$

Proof. First of all, we must see how to normalize the sequence of polygons Π_n so that they stay in a bounded region of the plane as $n \to \infty$. In our previous article we observed that the smallest square with sides parallel to the two axes Ox, Oy and containing Π_n, has sides of size $2P(n+1)-1$ where the recurrence

$$P(0) = 0, P(1) = 1; P(n) = 2P(n-1) + P(n-2), \quad \text{for } n > 1,$$

defines the so-called Pell numbers. Then they satisfy the relations

$$\begin{aligned} P(n+1) &= \frac{2+\sqrt{2}}{4}\left(1+\sqrt{2}\right)^n + \frac{2-\sqrt{2}}{4}\left(1-\sqrt{2}\right)^n \\ 2P(n+1)-1 &= \frac{2+\sqrt{2}}{2}\left(1+\sqrt{2}\right)^n + \frac{2-\sqrt{2}}{2}\left(1-\sqrt{2}\right)^n - 1 \\ &\sim \frac{2+\sqrt{2}}{2}\left(1+\sqrt{2}\right)^n. \end{aligned}$$

The polygon Π_n is composed of $4|q_{3n+1}|$ unit segments and blows up as $n \to \infty$. But the *normalized* polygon $\frac{1}{2P(n+1)-1}\Pi_n$ stays bounded. It has $4|q_{3n+1}|$ sides each of length $(2P(n+1)-1)^{-1}$. The total d-dimensional normalized polygon has length

$$\frac{4|q_{3n+1}|}{(2P(n+1)-1)^d}$$

and therefore the fractal dimension of the Fibonacci snowflake is

$$d = \lim_{n\to\infty} \frac{\log(|q_{3n+1}|)}{\log(P(n+1))} = \lim_{n\to\infty} \frac{\log\left(\frac{1+\sqrt{5}}{2}\right)^{3n+1}}{\log(1+\sqrt{2})^n} = \frac{\log(2+\sqrt{5})}{\log(1+\sqrt{2})}. \quad \clubsuit$$

§3. A measure of complexity. There are obviously many ways to measure the complexity of a line. We mention below a measure related to the number of points of intersection of the figure with a *random* straight line.

Recall a classical result due to Cauchy [2], Crofton [3], Steinhaus [5]. Consider a plane rectifiable curve Γ of length $|\Gamma|$ and whose convex hull K has a frontier of length $|\partial K|$. A straight line D is defined by

$$x\cos\theta + y\sin\theta - \rho = 0.$$

The probability measure considered is the uniform Lebesgue measure $d\rho.d\theta$ conditioned by the fact that D intersects Γ. The result we alluded to is that the average number of intersecting points of Γ with a random D is

$$N = \frac{2|\Gamma|}{|\partial K|}.$$

Applying this to Π_n, we see that the average number N_n of intersection points of Π_n with D is

$$N_n = 2\frac{2|q_{3n+1}|}{|\partial K_n|},$$

where K_n is the convex hull of Π_n. Easily seen, $|\partial K_n|$ is of the order of $P(n+1)$. Therefore,

Theorem 2. *We have*

$$N_n \sim a\frac{(2+\sqrt{5})^n}{(1+\sqrt{2})^n} = a\left(\frac{2+\sqrt{5}}{1+\sqrt{2}}\right)^n.$$

There is no difficulty to compute the constant $a = (1+\sqrt{5})$, but the important point is that the average number N_n of intersection points increases exponentially, showing once more the high complexity of the Fibonacci snowflake. Theorem 2 could obviously be considered as a corollary of the proof of Theorem 1. Both results are strongly related.

Remark 1. Let δ_n be the diameter of Π_n (which is of the order of $P(n+1)$). Obviously, $2\delta_n \leq |\partial K_n| \leq \pi\delta_n$. Theorem 2 states that the ratio $\mathcal{L}_n/\delta_n$ is of the order of N_n where $\mathcal{L}_n$ is the length of Π_n. The difference

$$\mathcal{L}_n - \delta_n = \delta_n \left(\frac{\mathcal{L}_n}{\delta_n} - 1 \right)$$

measures the distance of Π_n from being a straight line. The larger the ratio, more the curve meanders.

Remark 2. Let F be a positive strictly increasing function. The ratio $F(\mathcal{L}_n)/F(\delta_n)$ is a measure of the complexity of Π_n. When $F = \log$ we obtain Theorem 1 and when $F = \mathrm{Id}$ we obtain Theorem 2. The choice $F = \exp$ leads to $\exp(\mathcal{L}_n - \delta_n)$ i.e. Remark 1.

§4. Entropy. Let $p_j^{(n)}$ be the probability that a random straight line intersects Π_n in exactly j points, given that the line meets Π_n. The associated entropy h_n is by definition

$$h_n = -\sum p_j(n) \log(p_j(n)).$$

In [4] it is shown that

$$h_n \leq \log\left(\frac{2\mathcal{L}_n}{|\partial K_n|}\right) + \left(1 - \frac{|\partial K_n|}{2\mathcal{L}_n}\right) \log\left(\frac{2\mathcal{L}_n}{2\mathcal{L}_n - |\partial K_n|}\right).$$

The second term in the righthand side is positive, less than 1 and tends to 0 with $\frac{|\partial K_n|}{\mathcal{L}_n}$.

Since $\mathcal{L}_n$ is of the order of $(2+\sqrt{5})^n$ and $|\partial K_n|$ is of the order of $(1+\sqrt{2})^n$, we have

$$h_n \leq n \log\left(\frac{2+\sqrt{5}}{1+\sqrt{2}}\right) + \mathcal{O}(1).$$

We have thus established an upper bound for the complexity of Π_n:

Theorem 3.

$$\limsup_{n\to\infty} \frac{1}{n} h_n \leq \log\left(\frac{2+\sqrt{5}}{1+\sqrt{2}}\right).$$

References

[1] A. Blondin Massé, S. Brlek, S. Labbé, and M. Mendès France, *Fibonacci snowflakes*, Ann. Sci. Math. Québec 35 (2011), to appear.

[2] A.-L. Cauchy, *Mémoire sur la rectification des courbes et la quadrature des surfaces courbes*, Mém. Acad. Sci. 22 (1850), 3-15.

[3] M. W. Crofton, *On the theory of local probability, applied to straight lines drawn at random in a plane; the methods used being also extended to the proof of certain new theorems in the integral calculus*, Phil. Trans. of the royal soc. London 158 (1868), 181-199.

[4] M. Mendès France, *The Planck constant of a curve*, in Fractal Geometry and Analysis (J. Bélair and S. Dubuc, eds.), vol. 346, Kluwer Academic Publishers, 1991, pp. 325-366.

[5] H. Steinhaus, *Length, shape and area*, Colloq. Math. 3 (1954), 1-13.

A. Blondin Massé, S. Brlek, S. Labbé, LaCIM, Université du Québec à Montréal, C.P. 8888, Succ. Centre-ville, Montréal (QC) Canada H3C 3P8

E-mail address: **{blondin_masse.alexandre, labbe.sebastien}@courrier.uqam.ca, brlek.srecko@uqam.ca**

M. Mendès France, Département de mathématiques, UMR 5251, Université Bordeaux 1, 351 cours de la Libération, F-33405 Talence cedex, France

E-mail address: **michel.mendes-france@math.u-bordeaux1.fr**